

COVID-19 Vaccine Eligibility

Use this chart to help determine when to give COVID-19 vaccine based on age or health conditions.

Dose		Age:	12-17	18-49	50-64	65+
Pfizer	Primary 2 doses		✓	✓	✓	✓
	Additional (3rd) dose for immunocompromised at least 28 days after 2nd Pfizer dose		✓	✓	✓	✓
	Booster dose of Pfizer or different brand, at least 6 months after 2nd dose of Pfizer					✓
	• Long-term care facility resident			✓	✓	
	• Underlying medical condition or increased risk of social inequities			✓ may receive	✓	
	• Increased risk of exposure from work or institutional setting			✓ may receive	✓ may receive	
Moderna	Primary 2 doses			✓	✓	✓
	Additional (3rd) dose for immunocompromised at least 28 days after 2nd Moderna dose			✓	✓	✓
	Booster dose of Moderna (half-dose) or different brand, at least 6 months after 2nd dose of Moderna					✓
	• Long-term care facility resident			✓	✓	
	• Underlying medical condition or increased risk of social inequities			✓ may receive	✓	
	• Increased risk of exposure from work or institutional setting			✓ may receive	✓ may receive	
Johnson & Johnson	Primary 1 dose			✓	✓	✓
	Booster dose of J&J or different brand, at least 2 months after primary J&J dose			✓	✓	✓

View Related Guidance and Requirements for more information.